

A NEW OBSTRUCTION TO MINIMAL ISOMETRIC IMMERSIONS INTO A REAL SPACE FORM

TEODOR OPREA

University of Bucharest,
Faculty of Mathematics and Informatics
14 Academiei St., code 010014, Bucharest, Romania
e-mail: teodoroprea@yahoo.com

Abstract

In the theory of minimal submanifolds, the following problem is fundamental: *when does a given Riemannian manifold admit (or does not admit) a minimal isometric immersion into an Euclidean space of arbitrary dimension?* S.S. Chern, in his monograph *Minimal submanifolds in a Riemannian manifold*, remarked that the result of Takahashi (*the Ricci tensor of a minimal submanifold into a Euclidean space is negative semidefinite*) was the only known Riemannian obstruction to minimal isometric immersions in Euclidean spaces. A second solution to this problem was obtained by B.Y. Chen as an immediate application of his fundamental inequality [1]: *the scalar curvature and the sectional curvature of a minimal submanifold into a Euclidean space satisfies the inequality $\tau \leq k$* . In this paper we prove that the sectional curvature of a minimal submanifold into a Euclidean space also satisfies the inequality $k \leq -\tau$.

2000 Mathematics Subject Classification: 53C21, 53C24, 49K35.

Key words: constrained maximum, Chen's inequality, minimal submanifolds

1. OPTIMIZATIONS ON RIEMANNIAN MANIFOLDS

Let $(N, \tilde{g})$ be a Riemannian manifold, (M, g) a Riemannian submanifold, and $f \in \mathcal{F}(N)$. To these ingredients we attach the optimum problem

$$(1) \min_{x \in M} f(x).$$

Let's remember the result obtained in [6].

THEOREM 1.1. *If $x_0 \in M$ is the solution of the problem (1), then*

i) $(\text{grad } f)(x_0) \in T_{x_0}^\perp M,$

ii) *the bilinear form*

$$\begin{aligned} \alpha : T_{x_0} M \times T_{x_0} M &\rightarrow R, \\ \alpha(X, Y) &= \text{Hess}_f(X, Y) + \tilde{g}(h(X, Y), (\text{grad } f)(x_0)) \end{aligned}$$

is positive semidefinite, where h is the second fundamental form of the submanifold M in N .

2. THE CHEN'S INEQUALITY

Let (M, g) be a Riemannian manifold of dimension n , and x a point in M . We consider the orthonormal frame $\{e_1, e_2, \dots, e_n\}$ in $T_x M$.

The scalar curvature at x is defined by

$$\tau = \sum_{1 \leq i < j \leq n} R(e_i, e_j, e_i, e_j).$$

We denote

$$\delta_M = \tau - \min(k),$$

where k is the sectional curvature at the point x . The invariant δ_M is called the Chen's invariant of Riemannian manifold (M, g) .

The Chen's invariant can be estimated, if (M, g) is a Riemannian submanifold in a real space form $\widetilde{M}(c)$, varying with c and the mean curvature of M in $\widetilde{M}(c)$.

THEOREM 2.1. *Consider $(\widetilde{M}(c), \widetilde{g})$ a real space form of dimension m , $M \subset \widetilde{M}(c)$ a Riemannian submanifold of dimension $n \geq 3$. The Chen's invariant of M satisfies*

$$\delta_M \leq \frac{n-2}{2} \left\{ \frac{n^2}{n-1} \|H\|^2 + (n+1)c \right\},$$

where H is the mean curvature vector of submanifold M in $\widetilde{M}(c)$. The equality is attained at the point $x \in M$ if and only if there is an orthonormal frame $\{e_1, \dots, e_n\}$ in $T_x M$ and an orthonormal frame $\{e_{n+1}, \dots, e_m\}$ in $T_x^\perp M$ in which the Weingarten operators have the following form

$$A_{n+1} = \begin{pmatrix} h_{11}^{n+1} & 0 & 0 & \cdot & 0 \\ 0 & h_{22}^{n+1} & 0 & \cdot & 0 \\ 0 & 0 & h_{33}^{n+1} & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & h_{nn}^{n+1} \end{pmatrix},$$

with $h_{11}^{n+1} + h_{22}^{n+1} = h_{33}^{n+1} = \dots = h_{nn}^{n+1}$ and

$$A_r = \begin{pmatrix} h_{11}^r & h_{12}^r & 0 & \cdot & 0 \\ h_{12}^r & -h_{11}^r & 0 & \cdot & 0 \\ 0 & 0 & 0 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & 0 \end{pmatrix}, \quad r \in \overline{n+2, m}.$$

COROLLARY 2.1. *If the Riemannian manifold (M, g) , of dimension $n \geq 3$, admit a minimal isometric immersion into a real space form $\widetilde{M}(c)$, then*

$$k \geq \tau - \frac{(n-2)(n+1)c}{2}.$$

By using the constrained extremum method we give another Riemannian obstruction to minimal isometric immersions in real space forms

$$k \leq -\tau + \frac{(n^2 - n + 2)c}{2}.$$

3. A NEW OBSTRUCTION TO MINIMAL ISOMETRIC IMMERSIONS INTO A REAL SPACE FORM

Let (M, g) be a Riemannian manifold of dimension n . We define the following invariants

$$\delta_M^a = \begin{cases} \tau - a \min k, & \text{for } 0 \leq a < 1, \\ \tau - a \max k, & \text{for } -1 < a < 0, \end{cases}$$

$$\delta'_M = \tau + \max k,$$

where τ is the scalar curvature, and k is the sectional curvature.

With these ingredients we obtain

THEOREM 3.1. *For any real number $a \in (-1, 1)$, the invariant δ_M^a of a Riemannian submanifold (M, g) , of dimension $n \geq 3$, into a real space form $\widetilde{M}(c)$, of dimension m , verifies the inequality*

$$\delta_M^a \leq \frac{(n^2 - n - 2a)c}{2} + \frac{n(a+1) - 3a - 1}{n(a+1) - 2a} \frac{n^2 \|H\|^2}{2},$$

where H is the mean curvature vector of submanifold M in $\widetilde{M}(c)$. The equality is attained at the point $x \in M$ if and only if there is an orthonormal frame $\{e_1, \dots, e_n\}$ in $T_x M$ and an orthonormal frame $\{e_{n+1}, \dots, e_m\}$ in $T_x^\perp M$ in which the Weingarten operators have the following form

$$A_r = \begin{pmatrix} h_{11}^r & 0 & 0 & \cdot & 0 \\ 0 & h_{22}^r & 0 & \cdot & 0 \\ 0 & 0 & h_{33}^r & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & h_{nn}^r \end{pmatrix},$$

with $(a+1)h_{11}^r = (a+1)h_{22}^r = h_{33}^r = \dots = h_{nn}^r, \forall r \in \overline{n+1, m}$.

Proof. Consider $x \in M$, $\{e_1, e_2, \dots, e_n\}$ an orthonormal frame in $T_x M$, $\{e_{n+1}, e_{n+2}, \dots, e_m\}$ an orthonormal frame in $T_x^\perp M$ and $a \in (-1, 1)$.

From Gauss equation it follows

$$\begin{aligned} \tau - ak(e_1 \wedge e_2) &= \frac{(n^2-n-2a)c}{2} + \sum_{r=n+1}^m \sum_{1 \leq i < j \leq n} (h_{ii}^r h_{jj}^r - (h_{ij}^r)^2) - \\ &\quad - a \sum_{r=n+1}^m (h_{11}^r h_{22}^r - (h_{12}^r)^2). \end{aligned}$$

Using the fact that $a \in (-1, 1)$, we obtain

$$(1) \quad \tau - ak(e_1 \wedge e_2) \leq \frac{(n^2-n-2a)c}{2} + \sum_{r=n+1}^m \sum_{1 \leq i < j \leq n} h_{ii}^r h_{jj}^r - a \sum_{r=n+1}^m h_{11}^r h_{22}^r.$$

For $r \in \overline{n+1, m}$, let us consider the quadratic form

$$\begin{aligned} f_r : R^n &\rightarrow R, \\ f_r(h_{11}^r, h_{22}^r, \dots, h_{nn}^r) &= \sum_{1 \leq i < j \leq n} (h_{ii}^r h_{jj}^r) - ah_{11}^r h_{22}^r \end{aligned}$$

and the constrained extremum problem

$$\max f_r,$$

$$\text{subject to } P : h_{11}^r + h_{22}^r + \dots + h_{nn}^r = k^r,$$

where k^r is a real constant.

The first three partial derivatives of the function f_r are

$$\begin{aligned} (2) \quad \frac{\partial f_r}{\partial h_{11}^r} &= \sum_{2 \leq j \leq n} h_{jj}^r - ah_{22}^r, \\ (3) \quad \frac{\partial f_r}{\partial h_{22}^r} &= \sum_{j \in \overline{1, n} \setminus \{2\}} h_{jj}^r - ah_{11}^r, \\ (4) \quad \frac{\partial f_r}{\partial h_{33}^r} &= \sum_{j \in \overline{1, n} \setminus \{3\}} h_{jj}^r. \end{aligned}$$

As for a solution $(h_{11}^r, h_{22}^r, \dots, h_{nn}^r)$ of the problem in question, the vector $(\text{grad}) (f_1)$ is normal at P , from (2) and (3) we obtain $\sum_{j=1}^n h_{jj}^r - h_{11}^r -$

$$-ah_{22}^r = \sum_{j=1}^n h_{jj}^r - h_{22}^r - ah_{11}^r, \text{ therefore}$$

$$(5) \quad h_{11}^r = h_{22}^r = b^r.$$

- From (2) and (4), it follows $\sum_{j=1}^n h_{jj}^r - h_{11}^r - ah_{22}^r = \sum_{j=1}^n h_{jj}^r - h_{33}^r$. By using
- (5) we obtain $h_{33}^r = b^r(a+1)$. Similarly one gets
- (6) $h_{jj}^r = b^r(a+1), \forall j \in \overline{3, n}$.
- As $h_{11}^r + h_{22}^r + \dots + h_{nn}^r = k^r$, from (5) and (6) we obtain
- (7) $b^r = \frac{k^r}{n(a+1)-2a}$.

We fix an arbitrary point $p \in P$.

The 2-form $\alpha : T_p P \times T_p P \rightarrow R$ has the expression

$$\alpha(X, Y) = \text{Hess}_{f_r}(X, Y) + \langle h'(X, Y), (\text{grad } f_r)(p) \rangle,$$

where h' is the second fundamental form of P in R^n and $\langle \cdot, \cdot \rangle$ is the standard inner-product on R^n .

In the standard frame of R^n , the Hessian of f_r has the matrix

$$\text{Hess}_{f_r} = \begin{pmatrix} 0 & 1-a & 1 & \cdot & 1 \\ 1-a & 0 & 1 & \cdot & 1 \\ 1 & 1 & 0 & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & 1 & 1 & \cdot & 0 \end{pmatrix}.$$

As P is totally geodesic in R^n , considering a vector X tangent to P at the arbitrary point p , that is, verifying the relation $\sum_{i=1}^n X^i = 0$, we have

$$\begin{aligned} \alpha(X, X) &= 2 \sum_{1 \leq i < j \leq n} X^i X^j - 2aX^1 X^2 = \left(\sum_{i=1}^n X^i \right)^2 - \sum_{i=1}^n (X^i)^2 - 2aX^1 X^2 = \\ &= - \sum_{i=1}^n (X^i)^2 - a(X^1 + X^2)^2 + a(X^1)^2 + a(X^2)^2 = \\ &= - \sum_{i=3}^n (X^i)^2 - a(X^1 + X^2)^2 - (1-a)(X^1)^2 - (1-a)(X^2)^2 \leq 0. \end{aligned}$$

Therefore the point $(h_{11}^r, h_{22}^r, \dots, h_{nn}^r)$, which satisfies (5), (6), (7) is a maximum point.

From (5) and (6) it follows

$$\begin{aligned} (8) \quad f_r &\leq (b^r)^2 + 2b^r(n-2)b^r(a+1) + C_{n-2}^2(b^r)^2(a+1)^2 - a(b^r)^2 = \\ &= \frac{(b^r)^2}{2} [n^2(a+1)^2 - n(a+1)(5a+1) + 6a^2 + 2a] = \end{aligned}$$

$$= \frac{(b^r)^2}{2} [n(a+1) - 3a - 1][n(a+1) - 2a].$$

By using (7) and (8), we obtain

$$(9) \quad f_r \leq \frac{(k^r)^2}{2[n(a+1)-2a]} [n(a+1) - 3a - 1] = \frac{n^2(H^r)^2}{2} \frac{n(a+1)-3a-1}{n(a+1)-2a}.$$

The relations (1) and (9) imply

$$(10) \quad \tau - ak(e_1 \wedge e_2) \leq \frac{(n^2-n-2a)c}{2} + \frac{n(a+1)-3a-1}{n(a+1)-2a} \frac{n^2\|H\|^2}{2}.$$

In (10) we have equality if and only if the same thing occurs in the inequality (1) and, in addition, (5) and (6) occurs. Therefore

$$(11) \quad h_{ij}^r = 0, \quad \forall r \in \overline{n+1, m}, \quad \forall i, j \in \overline{1, n}, \quad \text{with } i \neq j \text{ and}$$

$$(12) \quad (a+1)h_{11}^r = (a+1)h_{22}^r = h_{33}^r = \dots = h_{nn}^r, \quad \forall r \in \overline{n+1, m}.$$

The relations (10), (11) and (12) imply the conclusion of the theorem.

Remark. i) Making a to converge at 1 in previous inequality, we obtain *Chen's Inequality*. The conditions for which we have equality are obtained in [1] and [6].

ii) For $a = 0$ we obtain the well-known inequality

$$\tau \leq \frac{n(n-1)}{2} (\|H\|^2 + c).$$

The equality is attained at the point $x \in M$ if and only if x is a totally umbilical point.

iii) Making a to converge at -1 in previous inequality, we obtain

$$\delta'_M \leq \frac{(n^2 - n + 2)c}{2} + \frac{n^2 \|H\|^2}{2}.$$

THEOREM 3.2. *The invariant δ'_M of a Riemannian submanifold (M, g) , of dimension $n \geq 3$, into a real space form $\widetilde{M}(c)$, of dimension m , verifies the inequality*

$$\delta'_M \leq \frac{(n^2 - n + 2)c}{2} + \frac{n^2 \|H\|^2}{2},$$

where H is the mean curvature vector of submanifold M in $\widetilde{M}(c)$. The equality is attained at the point $x \in M$ if and only if there is an orthonormal

frame $\{e_1, \dots, e_n\}$ in $T_x M$ and an orthonormal frame $\{e_{n+1}, \dots, e_m\}$ in $T_x^\perp M$ in which the Weingarten operators have the following form

$$A_r = \begin{pmatrix} h_{11}^r & 0 & 0 & \cdot & 0 \\ 0 & h_{22}^r & 0 & \cdot & 0 \\ 0 & 0 & 0 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & 0 \end{pmatrix},$$

with $h_{11}^r = h_{22}^r, \forall r \in \overline{n+1, m}$.

Proof. We consider the point $x \in M$, the orthonormal frames $\{e_1, \dots, e_n\}$ in $T_x M$ and $\{e_{n+1}, \dots, e_m\}$ in $T_x^\perp M$, $\{e_1, e_2\}$ being an orthonormal frame in the 2- plane which maximize the sectional curvature at the point x in $T_x M$.

The invariant δ'_M verifies

$$\begin{aligned} (1) \quad \delta'_M &= \frac{(n^2-n+2)c}{2} + \sum_{r=n+1}^m \sum_{1 \leq i < j \leq n} (h_{ii}^r h_{jj}^r - (h_{ij}^r)^2) + \sum_{r=n+1}^m (h_{11}^r h_{22}^r - (h_{12}^r)^2) \leq \\ &\leq \frac{(n^2-n+2)c}{2} + \sum_{r=n+1}^m \sum_{1 \leq i < j \leq n} (h_{ii}^r h_{jj}^r) + h_{11}^r h_{22}^r. \end{aligned}$$

For $r \in \overline{n+1, m}$, let us consider the quadratic form

$$\begin{aligned} f_r : R^n &\rightarrow R, \\ f_r(h_{11}^r, h_{22}^r, \dots, h_{nn}^r) &= \sum_{1 \leq i < j \leq n} (h_{ii}^r h_{jj}^r) + h_{11}^r h_{22}^r \end{aligned}$$

and the constrained extremum problem

$$\max f_r,$$

$$\text{subject to } P : h_{11}^r + h_{22}^r + \dots + h_{nn}^r = k^r,$$

where k^r is a real constant.

The first three partial derivatives of the function f_r are

$$\begin{aligned} (2) \quad \frac{\partial f_r}{\partial h_{11}^r} &= \sum_{2 \leq j \leq n} h_{jj}^r + h_{22}^r, \\ (3) \quad \frac{\partial f_r}{\partial h_{22}^r} &= \sum_{j \in \overline{1, n} \setminus \{2\}} h_{jj}^r + h_{11}^r, \\ (4) \quad \frac{\partial f_r}{\partial h_{33}^r} &= \sum_{j \in \overline{1, n} \setminus \{3\}} h_{jj}^r. \end{aligned}$$

As for a solution $(h_{11}^r, h_{22}^r, \dots, h_{nn}^r)$ of the problem in question, the vector $(\text{grad}) (f_1)$ is normal at P , from (2) and (3) we obtain $\sum_{j=1}^n h_{jj}^r - h_{11}^r +$
 $+h_{22}^r = \sum_{j=1}^n h_{jj}^r - h_{22}^r + h_{11}^r$, therefore
(5) $h_{11}^r = h_{22}^r = b^r$.

From (2) and (4), it follows $\sum_{j=1}^n h_{jj}^r - h_{11}^r + h_{22}^r = \sum_{j=1}^n h_{jj}^r - h_{33}^r$. By using
(5) we obtain $h_{33}^r = 0$. Similarly one gets
(6) $h_{jj}^r = 0, \forall j \in \overline{3, n}$.

As $h_{11}^r + h_{22}^r + \dots + h_{nn}^r = k^r$, from (5) and (6) we obtain
(7) $b^r = \frac{k^r}{2}$.

We fix an arbitrary point $p \in P$.

The 2-form $\alpha : T_p P \times T_p P \rightarrow R$ has the expression

$$\alpha(X, Y) = \text{Hess}_{f_r}(X, Y) + \langle h'(X, Y), (\text{grad } f_r)(p) \rangle,$$

where h' is the second fundamental form of P in R^n and $\langle \cdot, \cdot \rangle$ is the standard inner-product on R^n .

In the standard frame of R^n , the Hessian of f_r has the matrix

$$\text{Hess}_{f_r} = \begin{pmatrix} 0 & 2 & 1 & \cdot & 1 \\ 2 & 0 & 1 & \cdot & 1 \\ 1 & 1 & 0 & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & 1 & 1 & \cdot & 0 \end{pmatrix}.$$

As P is totally geodesic in R^n , considering a vector X tangent to P at the arbitrary point p , that is, verifying the relation $\sum_{i=1}^n X^i = 0$, we have $\alpha(X, X) = 2 \sum_{1 \leq i < j \leq n} X^i X^j + 2X^1 X^2 = (\sum_{i=1}^n X^i)^2 - \sum_{i=1}^n (X^i)^2 + 2X^1 X^2 = - \sum_{i=3}^n (X^i)^2 - (X^1 - X^2)^2$.

The 2-form α is semipositive definite. Therefore the point $(h_{11}^r, h_{22}^r, \dots, h_{nn}^r)$ which satisfies (5), (6) and (7) is a global maximum point. Using this fact and (5), (6) and (7) we obtain

$$\delta'_M \leq \frac{(n^2 - n + 2)c}{2} + \frac{n^2 \|H\|^2}{2}$$

The relation

$$\delta'_M = \frac{(n^2 - n + 2)c}{2} + \frac{n^2 \|H\|^2}{2}$$

occurs if and only if we have

- (9) $h_{ij}^r = 0, \forall r \in \overline{n+1, m}, \forall i, j \in \overline{1, n}$ with $i \neq j$,
- (10) $h_{11}^r = h_{22}^r, \forall r \in \overline{n+1, m}$,
- (11) $h_{33}^r = \dots = h_{nn}^r = 0, \forall r \in \overline{n+1, m}$.

Therefore, there is an orthonormal frame $\{e_1, \dots, e_n\}$ in $T_x M$ and an orthonormal frame $\{e_{n+1}, \dots, e_m\}$ in $T_x^\perp M$ in which the Weingarten operators have the following form

$$A_r = \begin{pmatrix} h_{11}^r & 0 & 0 & \cdot & 0 \\ 0 & h_{22}^r & 0 & \cdot & 0 \\ 0 & 0 & 0 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & 0 \end{pmatrix},$$

with $h_{11}^r = h_{22}^r, \forall r \in \overline{n+1, m}$.

COROLLARY 3.1. *If the Riemannian manifold (M, g) , of dimension $n \geq 3$, admit a minimal isometric immersion into a real space form $\widetilde{M}(c)$, then*

$$\tau - \frac{(n-2)(n+1)c}{2} \leq k \leq -\tau + \frac{(n^2 - n + 2)c}{2}.$$

COROLLARY 3.2. *If the Riemannian manifold (M, g) , of dimension $n \geq 3$, admit a minimal isometric immersion into a Euclidean space, then*

$$\tau \leq k \leq -\tau.$$

References

- [1] B.Y. Chen, *Some pinching classification theorems for minimal submanifolds*. Arch. Math. **60** (1993), 568-578.
- [2] B.Y. Chen, *A Riemannian invariant for submanifolds in space forms its applications*. Geom. Topology of submanifolds, World Scientific, Leuven, Brussel **VI** (1993), 58-81.
- [3] B.Y. Chen, *A Riemannian invariant its applications to submanifolds theory*. Results in Mathematics **27** (1995), 17-26.
- [4] B.Y. Chen, *Mean curvature and shape operator of isometric immersions in real-space-forms*. Glasgow Math. J. **38** (1996), 87-97.
- [5] B.Y. Chen, *Some new obstructions to minimal Lagrangian isometric immersions*. Japan. J. Math. **26** (2000), 105-127.
- [6] T. Oprea, *Optimizations on Riemannian Submanifolds*. An. Univ. Buc., **LIV**, 1 (2005), 127-136.
- [7] C. Udriște, *Convex functions and optimization methods on Riemannian manifolds*. Mathematics its Applications, **297**, Kluwer Academic Publishers Group, Dordrecht, 1994.
- [8] C. Udriște, O. Dogaru and I. Tevy, *Extrema with nonholonomic constraints*. Geometry Balkan Press, Bucharest, 2002.